



Practical target strength of free-swimming chub mackerel *Scomber japonicus*

Yanhui Zhu¹ · Keizo Ito² · Kosuke Mizutani² · Kenji Minami¹ · Hokuto Shirakawa³ · Yohei Kawauchi⁴ · Yuka Iwahara⁵ · Kimiharu Nahata² · Nobuhiko Sato¹ · Kyosuke Seki² · Mitsuki Kuroda² · Kazushi Miyashita¹

Received: 13 April 2023 / Accepted: 25 October 2023

© The Author(s), under exclusive licence to Japanese Society of Fisheries Science 2023

Abstract

Chub mackerel *Scomber japonicus* is a migratory fish widely distributed around Japan, and is an important fishery resource. However, target strength (*TS*) measurements of chub mackerel are limited, and the relationship between *TS* and fork length has not been fully clarified, despite its importance for the estimation of chub mackerel abundance. In this study, the *TS*–fork length (*FL*) relationship in chub mackerel was evaluated under realistic conditions. TS_{mean} and TS_{max} tended to increase with fork length at both 38 and 120 kHz, and *TS* histograms were bimodal for most individuals. In the TS_{mean} –*FL* relationship, when the coefficient *a* was fixed at 20 (the standard for fish with swim bladders), TS_{cm} (standardized by the square of the fork length) was -67.9 dB ($r^2=0.70$) at 38 kHz and -69.2 dB ($r^2=0.45$) at 120 kHz. Additionally, the swimming angle had a peak of around 0° with a mean of -1.23° , and the mean swimming speed was 0.16 *FL*/s with a standard deviation of 0.07 *FL*/s. *TS* reached a maximum between -20° and 0° and then decreased drastically as the swimming angle increased or decreased. Our results have practical implications for the management of chub mackerel.

Keywords Chub mackerel · Target strength · Free-swimming · Swimming behavior · Quantitative echo sounder

✉ Yanhui Zhu
zhuyanhui920817@gmail.com

Keizo Ito
itoh.keizo.i8@elms.hokudai.ac.jp

Kosuke Mizutani
go.oden0301@gmail.com

Kenji Minami
k.minami@fsc.hokudai.ac.jp

Hokuto Shirakawa
shirakawa_hokuto26@fra.go.jp

Yohei Kawauchi
kawauchi_yohei41@fra.go.jp

Yuka Iwahara
iwahara_yuka58@fra.go.jp

Kimiharu Nahata
kimi1230otaru@eis.hokudai.ac.jp

Nobuhiko Sato
sato.n.nobuhiko@gmail.com

Kyosuke Seki
seki.kyosuke.i1@elms.hokudai.ac.jp

Mitsuki Kuroda
mitsuki_k_fishing@eis.hokudai.ac.jp

Kazushi Miyashita
miyashi@fish.hokudai.ac.jp

¹ Field Science Center for Northern Biosphere, Hokkaido University, 20-5 Benten-cho, Hakodate, Hokkaido 040-0051, Japan

² Graduate School of Environmental Science, Hokkaido University, 20-5 Benten-cho, Hakodate, Hokkaido 040-0051, Japan

³ Demersal Fish Resources Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, National Research and Development Agency, Japan Fisheries Research and Education Agency, 1-5939-22, Suido-cho, Chuo-ku, Niigata-shi, Niigata 951-8121, Japan

⁴ Socio-Ecological Systems Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 2-12-4 Fukuura, Kanazawa, Yokohama, Kanagawa 236-8648, Japan

⁵ Fisheries Data Science Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, Japan Fisheries Research and Education Agency, National Research and Development Agency, 2-12-4 Fukuura, Kanazawa, Yokohama, Kanagawa 236-8648, Japan

Introduction

The chub mackerel *Scomber japonicus* is a migratory fish widely distributed around Japan and an important fishery resource in Japan (Okada 2020). This species has been designated as a target for determination of total allowable catch (TAC) in Japan (Yatsu et al. 2019), and thus stock assessment and management have been conducted for the Pacific stock and the Tsushima Warm Current stock (Kurota et al. 2019). Chub mackerel is a small pelagic and mass-caught fish, and its abundance fluctuates substantially (Goto 2020). Fishing pressure is high for the species, and stock declines are likely to occur due to inadequate management, especially during periods of declining recruitment (Kawai et al. 2002; Yatsu and Kaeriyama 2005). In addition, since the distribution of chub mackerel depends on its abundance and environmental factors (Goto et al. 2020; Sassa and Tsukamoto 2010), it is important to clarify the characteristics of its distribution according to abundance levels (Kurota et al. 2019). For these reasons, detailed information on changes in abundance and fishing grounds of chub mackerel is essential for effective stock management.

Cohort analyses (e.g., virtual population analysis [VPA]) are the main approach for the assessment of chub mackerel in Japan (Nakayama and Hiramatsu 2010; Ichinokawa et al. 2017). In VPA, the spawning stock biomass and recruitment number are calculated based on the age-specific fishing mortality coefficient (Ichinokawa and Okamura 2014). Since the estimation recruitment accuracy using VPA increases with the number of years of data accumulation, it is difficult to grasp the abundance of age-stock at an early life stage (Shimura et al. 2009; Muko and Kurota 2018). In addition, the VPA assumes that the catch is proportional to abundance, and the estimation accuracy may be low for fish species with large fluctuations in abundance (Furusawa 1991). Thus, to estimate the abundance of chub mackerel more accurately, VPA needs to be corrected using recent resource indices (Ichinokawa and Okamura 2014). Therefore, there is a need for a method that enables the rapid and accurate estimation of the present abundance level and distribution.

Recently, acoustic monitoring methods that enable the quantitative visualization of fishery resources have attracted attention (Miyashita 2019; Munoz et al. 2020; Zhu et al. 2021a, b). Among these, a quantitative echo sounder (hereafter, echo sounder) can be applied to a wide area in a short time and can be used to estimate the abundance quickly and quantitatively, independent of the fishery catch (Abe et al. 1999; Zhu et al. 2021a, b). The volume backscatter strength (SV , dB) or area backscatter strength (SA , dB) obtained from the echo sounder is

obtained and the biological density is calculated using the target strength (TS , dB), indicating the strength per target organism, to estimate the abundance (Amakasu et al. 2003; Hirose et al. 2005; Tong et al. 2022). Therefore, the TS of target organisms is very important for accurate abundance estimation (Tong et al. 2022). However, the relationship between TS and fork length in chub mackerel has not been fully clarified owing to limited TS data (Miyanoohana et al. 1990; Mukai et al. 1994; Miyanoohana 1994). Thus, it is important to collect more TS data and obtain a TS -fork length equation with high accuracy for the chub mackerel stock assessment.

Since TS depends on the swimming posture of fish, measurements should be obtained under natural conditions (in situ method) (Sawada 2002). However, it is difficult to accurately determine the species and target fish length in the natural environment (Simmonds and MacLennan 2005). Therefore, measurements are frequently obtained in a controlled environment (i.e., ex situ method). The suspension and cage methods are common ex situ methods; however, these approaches restrict swimming behavior, and it is difficult to obtain practical TS values close to those in natural conditions (Mukai et al. 1994). To obtain robust and accurate TS data, it is important to evaluate target fish swimming freely in a physically controlled environment.

Therefore, the objective of this study was to accurately estimate the TS -fork length relationship by measuring the TS of free-swimming chub mackerel in a large indoor aquarium with a stable physical environment.

Materials and methods

TS measurement

Eighteen individuals (fork length [FL]: 17.4–34.0 cm) caught by angling along the coast of Hakodate, Hokkaido, during March 2021, August 2021, and December 2022, were used as experimental fish (Table 1). The caught fish were immediately transported to the Hakodate Research Center for Fisheries and Oceans (hereafter referred to as the Research Center). After acclimation, they were fed and reared in a 2500-L vinyl aquarium ($D \times H$: 2×0.8 m).

Two types of TS measurements were conducted in a large indoor aquarium ($W \times L \times H$: $5 \times 10 \times 6$ m) at the Research Center using two types of nets. Between March 8, 2021, and April 7, 2021, we used a circular net ($D \times H$: 3×6 m), which was placed in the center of the large aquarium (Fig. 1a). Between August 23 and August 27, 2021, and December 1 and December 13, 2022, we used a nylon partition net. The nylon partition net was installed from the water surface to the bottom of the aquarium, dividing the long side of the aquarium into two equal parts, and one half of the aquarium

Table 1 Experiment date and *TS* of each experimental individual

Date	Net Type	Fork length (<i>FL</i> , cm)	<i>TS</i> _{mean} (dB)		<i>TS</i> _{max} (dB)	
			38 kHz	120 kHz	38 kHz	120 kHz
8 March 2021 ~ 7 April 2021	Circular net	17.4	-43.9	-43.4	-36.1	-33.9
		24.1	-40.6	-47.3	-35.9	-33.8
		24.3	-39.1	-40.4	-34.6	-32.2
		28.0	-39.4	-40.9	-33.4	-31.1
23 August 2021 ~ 27 August 2021	Partition net	20.3	-39.9	-43.4	-34.4	-31.6
		22.0	-38.8	-41.5	-35.2	-31.6
		25.6	-37.2	-39.4	-32.5	-30.1
		34.0	-35.6	-39.9	-30.0	-30.0
1 December 2022 ~ 13 December 2022	Partition net	17.8	-42.8	-42.2	-37.8	-34.8
		17.9	-43.0	-46.0	-38.0	-35.0
		19.3	-41.9	-42.6	-36.4	-33.3
		19.4	-42.3	-46.7	-37.6	-34.8
		19.5	-44.9	-44.8	-37.4	-34.8
		21.5	-43.7	-41.5	-36.5	-32.3
		22.3	-42.5	-42.6	-36.7	-32.5
		22.4	-41.6	-41.1	-35.4	-32.3
		24.2	-40.6	-39.2	-35.9	-32.3
27.4	-39.5	-37.7	-32.5	-30.0		




Fig. 1 Experimental design. **a** Experiment with the circular net. **b** Experiment with the partition net

was used as the experimental area (Fig. 1b). For *TS* measurements, one individual was placed into the aquarium at a time.

In all experiments, a split-beam echo sounder (KSE-300, Sonic; frequencies: 38 and 120 kHz) capable of obtaining three-dimensional locations of fish was used (Table 2), and acoustic data were recorded for at least 12 h. During the experiment, the transducers of the echo sounder were fixed to a movable bridge, and the transducer surface was

positioned at about 40 cm below the water surface in the center of the experimental area. The positions of the transducer were fine-tuned so that the net did not interfere with the beam of the echo sounder. Before the start of each experiment, a tungsten carbide metal standard sphere (φ38.1 mm), which is a standard reflector with a known *TS*, was used to calibrate the echo sounder. Since air bubbles adhering to the gills of experimental fish can cause large measurement errors (Sawada 2002), efforts were made to prevent the fish

Table 2 Specification of quantitative echo sounder KSE-300 during *TS* measurements

Transducer	T-178	T-182
Frequency (kHz)	38	120
−3 dB beam width (degrees)	8.4	8.2
Transducer type		Split beam
Pulse length (ms)		0.3
Ping interval (s)		0.2

from encountering air during transport to the experimental aquarium. Furthermore, since water temperature influences the swimming behavior of fish (Wardle 1980), it must be controlled for *TS* measurements. The appropriate water temperature for chub mackerel varies depending on the season, with ranges of 14.1–16.4 °C from January to April and 17.7–21.7 °C from May to December (Okunishi et al. 2020). Therefore, the water temperature in the experimental aquarium was kept at 14 ± 1 °C during the March experiment and 18 ± 1 °C during the August and December experiment to prevent a decrease in the swimming ability of chub mackerel.

Analysis of acoustic data

TS values were determined by the algorithm for *TS* detection using the echo sounder and output as a CSV file (Sawada 2002). Since echoes from garbage in the water

can be misidentified as echoes of chub mackerel, unwanted echoes were excluded based on the signal-to-noise ratio (SNR). The background noise during the measurements in this study was approximately −80 dB or less, and a SNR of 20 dB or more is sufficient to accurately estimate *TS* (Sawada 2002); therefore, the lower threshold of *TS* was set to −60 dB. In addition, due to the performance of the echo sounder, accurate *TS* values cannot be obtained near the transducer and near the bottom (Kawauchi et al. 2019); accordingly, these data need to be excluded from the analysis. For the data near the transducers, the boundary of the far-field (*Rend*, m) (Medwin and Clay 1988), where sound pressure is stable and *TS* can be measured with high accuracy, and the diameter of the first Fresnel zone (*z*, m) (Yilmaz 1987), where the full length of the experimental fish must fit, were used as standards. Data shallower than that standard depth were excluded (Table 3). Next, data obtained up to 0.24 m from the bottom, corresponding

Table 3 Upper analytical depth based on the acoustic far-field boundary (*Rend*, m) and the diameter of the first Fresnel zone (*z*, m)

Fork length (<i>L</i> , cm)	<i>Rend</i> (m)		Measurable <i>L</i> under <i>Rend</i> (cm)			Final upper threshold (m)	
	38 kHz	120 kHz	38 kHz	120 kHz	120 kHz	38 kHz	120 kHz
17.4	1.55	0.62	35.0	12.5	1.21	1.55	1.21
17.8					1.27		1.27
17.9					1.28		1.28
19.3					1.49		1.49
19.4					1.51		1.51
19.5					1.52		1.52
20.3					1.65		1.65
21.5					1.85		1.85
22.0					1.94		1.94
22.3					1.99		1.99
22.4					2.01		2.01
24.1					2.32		2.32
24.2					2.34		2.34
24.3					2.36		2.36
25.6					2.62		2.62
27.4					3.00		3.00
28.0					3.14		3.14
34.0					4.62		4.62

to the acoustic blind zone, were excluded (Mitson 1983; Simmonds et al. 1992).

Using the TS data for chub mackerel extracted by the above procedure, the mean TS (TS_{mean}) and maximum TS (TS_{max}) values were obtained for each experimental condition. The equation for the TS – FL relationship was estimated by the least-squares method. In general, the TS – FL relationship is expressed by the following equation.

$$TS = a10\log_{10}FL + b \quad (1)$$

where a and b are coefficients and FL is the fork length (cm). It has been theoretically (Furusawa and Miyanoana 1988) and experimentally (Miyanoana et al. 1990) proven that the TS of fish with swim bladders, such as chub mackerel, is proportional to the square of the fork length. In this case, a is fixed at 20 and b is TS_{cm} standardized by the square of the fork length, which is expressed by the following equation.

$$TS = 20\log_{10}FL + TS_{\text{cm}} \quad (2)$$

At the time that TS data were recorded, changes in three-dimensional coordinate information as the fish swam were also measured for each ping. Using this coordinate information, it is possible to estimate the swimming speed and absolute swimming angle using the echo tracing method (Henderson et al. 2008; Furusawa and Amakasu 2010). The swimming angle obtained in this study is the angle of elevation/degradation in the direction the fish is moving, which is different from the exact posture angle. However, since the direction of the fish's body changes in accordance with the direction the fish is moving, it is possible to use the swimming angle as an indicator of the posture change. For this reason, the relationship between TS and swimming behavior of chub mackerel was examined from these swimming data. In this study, only tracking data for which echoes were taken for more than five pings in a row and for cases of straight swimming (i.e., low tortuosity: $Tor \leq 1.2$) (Henderson et al. 2008) were used.

The TS , swimming angle, and swimming speed were calculated in R ver. 4.1.0. Additionally, the parameters of TS – FL relationships were estimated using nonlinear least square methods by the nls function in R. Since tracking data were more obtained at 120 kHz than at 38 kHz, the swimming angle and swimming speed were verified by pooling data for all individuals at 120 kHz.

Results

For each individual, 13–9713 pings were obtained at 38 kHz, and 24–9577 pings were obtained at 120 kHz. Over 100 pings at both frequencies were obtained for more than 83% of individuals. The TS histograms were bimodal for most

individuals (Figs. 2 and 3). In addition, TS_{mean} values calculated from the measured data ranged from -44.9 to -35.6 dB at 38 kHz and from -47.3 to -37.7 dB at 120 kHz, while TS_{max} values ranged from -38.0 to -30.0 dB at 38 kHz and from -35.0 to -30.0 dB at 120 kHz (Table 1). At both frequencies, TS_{mean} and TS_{max} tended to increase with fork length. Furthermore, TS_{mean} was stronger at 38 kHz than at 120 kHz in most individuals. At TS_{max} , the pattern inverted, and values were stronger at 120 kHz than at 38 kHz.

The relationships between fork length and TS_{mean} , and between fork length and TS_{max} at both frequencies were obtained as shown in Fig. 4. In the relationship between fork length and TS_{mean} , when the coefficients were not fixed, a and b were 27.4 and -77.9 dB ($r^2=0.75$) at 38 kHz, respectively, and 23.3 and -73.7 dB ($r^2=0.46$) at 120 kHz, respectively. When the coefficient a was fixed at 20, TS_{cm} was -67.9 dB ($r^2=0.70$) at 38 kHz and -69.2 dB ($r^2=0.45$) at 120 kHz. In the relationship between fork length and TS_{max} , when the coefficients were not fixed, a and b were 24.9 and -68.9 dB ($r^2=0.96$) at 38 kHz and 20.1 and -59.6 dB ($r^2=0.96$) at 120 kHz, respectively. When the coefficient a was fixed at 20, TS_{cm} was -62.3 dB ($r^2=0.97$) at 38 kHz and -59.5 dB ($r^2=0.96$) at 120 kHz.

The swimming angle estimated in this study had a peak of around 0° , with a mean of -1.23° and a standard deviation of 11.50° (Fig. 5a). The mean swimming speed was 0.16 FL/s with a standard deviation of 0.07 FL/s (Fig. 5b). In addition, the relationship between the swimming angle and TS_{cm} , which was calculated using measured TS and FL , is shown in Fig. 6. In all individuals, TS_{cm} reached a maximum between -20° and 0° , and then decreased drastically as the swimming angle increased or decreased.

Discussion

Swimming behavior

Although the swimming angle is different from the posture angle, it is an indicator of postural changes that affect TS (Kawauchi et al. 2019). Here, we compared the swimming angle and swimming speed with those previously estimated for chub mackerel. Nauen et al. (2002) evaluated the swimming angle and posture of chub mackerel in an aquarium using a video camera, revealing that the fish tended to turn their bodies downward by approximately 3° relative to the direction of travel. Similar results were obtained by Gibb et al. (1999), who found that chub mackerel tended to swim with a downward tilt at all swimming speeds. In addition, Ito et al. (2015) tracked echoes of chub mackerel schools in the field using a broadband quantitative echo sounder with 80–120 kHz frequency on a research vessel, resulting a peak of slight smaller value than 0° for the swimming angle.




Fig. 2 TS histograms at 38 kHz. FL indicates the fork length of all individuals, and n indicates the number of TS detected by the echo sounder




Fig. 3 TS histograms at 120 kHz. FL indicates the fork length of all individuals, and n indicates the number of TS detected by the echo sounder




Fig. 4 Relationship between fork length (FL) and estimated TS_{mean} (a), and between fork length (FL) and estimated TS_{max} , (b) of chub mackerel in this study

Therefore, it seems that chub mackerels often swim horizontally or downward with a slight body tilt. In the present study, the estimated swimming angle had a peak of around 0° with a mean of -1.23° , indicating that the results of this study are consistent with those of previous studies on swimming behavior.

Additionally, in a previous study of the swimming speed of chub mackerel in the natural environment, as determined using a video camera in an underwater bulb, the average swimming speed of chub mackerel was $1.73 L/s$ (Hasegawa and Kobayashi 1989). However, the mean swimming speed in this study was $0.16 FL/s$, which was slower than


that reported by Hasegawa et al. (1989). This difference in swimming speed is thought to be caused by the difference in swimming between solitary animals and groups. In a study of rose bitterling *Rhodeus ocellatus ocellatus*, the swimming speed of the group increased as the number of individuals increased (Kanehiro et al. 1985). Therefore, our results could be interpreted to obtain a part of the main behavior.

As described above, the swimming angles obtained in this study were roughly similar to those measured in previous studies, suggesting that the behaviors observed in this study were not abnormal. However, the swimming speed of the chub mackerel in this study may not fully reflect the swimming behavior of chub mackerel in the wild. Therefore, it is important to improve the accuracy of estimation by measuring the TS of multiple individuals for fish species that swim in schools. In the future, by increasing the sample size, it will be possible to conduct measurements in swimming conditions similar to those in actual sea areas, leading to more accurate and practical TS estimation.

Relationship between the swimming angle and TS_{cm}

The TS_{cm} values in this study tended to be larger when fish were swimming slightly downward, with swimming angles ranging from -20° to 0° . This may be due to the increase in the projected area of the swim bladder in the dorsal direction during downward movement and the synchronous enhancement of acoustic reflection. In most fish with a swim bladder, the swim bladder is tilted upward relative to the fish body (Amakasu et al. 2010), and the dorsal cross-sectional area of the swim bladder is maximized when the fish head is tilted downward (Tanigami et al. 2006). Since the strongest acoustic reflector in the fish body is considered to be due to the swim bladder (Foote 1980), the TS tends to increase as the cross-sectional area of the swim bladder increases. Therefore, in this study, the peak TS_{cm} was observed when the chub mackerel swam slightly downward, which may reflect

Fig. 5 Histogram of a swimming angle and b swimming speed at 120 kHz; n indicates the number of swimming tracks






Fig. 6 Relationship between the swimming angle and TS_{cm} for experimental individuals at 120 kHz. Black circles indicate measured values

the upward orientation of the swim bladder in relation to the body axis.

Measured TS

In this study, chub mackerel showed a length-dependent trend in which TS increased with increasing fork length at both frequencies. In general, the TS of fish with a swim bladder is proportional to the square of the fork length. The slope a of the $TS_{mean}-FL$ relationship estimated in this study was 27.4 at 38 kHz and 23.3 at 120 kHz, which is close to 20. Additionally, the slope a of the $TS_{max}-FL$ equation was close to 20 at both frequencies. In general, when a was fixed at 20, the TS_{cm} of maximums TS ($Max\ TS_{cm}$) in the dorsal direction of fish with a swim bladder did not reflect the effect of posture angle, and $Max\ TS_{cm}$ calculated from a quantitative echo sounder was almost -59 to -61 dB, regardless of the frequency (Miyanohana 1994). In this study, the estimated $Max\ TS_{cm}$ values were -62.3 dB at 38 kHz and -59.5 dB at 120 kHz, which are close to the theoretical values. Therefore, the TS values measured in this study were considered reasonable.

In addition, the TS histograms of chub mackerel showed a bimodal distribution at both frequencies. Similar bimodal distributions have been obtained for related species and other taxa which have a similar type of swim bladder shape, including the Atlantic mackerel *Scomber scombrus* measured by in situ methods (Palermi et al. 2021) and the free-swimming Japanese jack mackerel *Trachurus japonicus* (Kawauchi et al. 2019). This distribution may be explained by the high directionality of TS with respect to posture angle (Miyanohana et al. 1990) and the sharp decrease in TS values with an increasing or decreasing posture angle from the peak

of the main lobe. With a wide range of swimming posture data, it is highly likely that TS histograms will be bimodal.

Furthermore, for many length classes, TS_{mean} was larger at 38 kHz than at 120 kHz; however, this pattern was reversed at TS_{max} , with greater variation in TS at higher frequencies. Similar results were obtained by Mukai et al. (1993). This is thought to be because directionality becomes stronger at high frequencies (Matsukura et al. 2013), and even small changes in posture angle cause large fluctuations in TS values. Additionally, in bladder fish, TS fluctuates significantly with posture when the standardized body length L/λ (i.e., the body length L standardized by wavelength λ) is large (Furusawa et al. 1989). In the case of chub mackerel in the length range covered in this study, L/λ was smaller at 38 kHz than at 120 kHz, and thus TS could be measured more stably at 38 kHz. These results suggest that low frequencies are suitable for future analyses of the abundance of adult chub mackerel, as estimates are less affected by the posture angle.

Equations relating TS_{mean} to FL

We compared the newly estimated $TS_{mean}-FL$ equations with those for the chub mackerel and related species reported in the literature (Table 4, Fig. 7). The TS_{cm} of chub mackerel (20.0–30.0 cm) estimated using in situ methods by Kuznetsov et al. (2021) were approximately 4 dB weaker than those of the present study at both frequencies. One of the reasons for this difference is that TS measurements obtained by in situ methods may include other fish species, since the target fish cannot be confirmed (Sawada 2002). The uncertainty in species identification might affect the accuracy of the TS , resulting the lower TS_{cm} in the previous studies.

On the other hand, the theoretically estimated TS_{cm} at 38 kHz was -66.0 dB for adult chub mackerel (16.0–28.0 cm), slightly different from 1.9 dB in the present study but 3.2 dB higher than that at 120 kHz in the present study (Park et al. 2022). Similar results for adult chub mackerel (26.2–38.3 cm) were found by Lee et al. (2005), where differences were small at low frequencies and became larger at high frequencies. This large difference at 120 kHz is thought to be due to a difference in the assumed mean of posture angles, and the change in TS with respect to the change in posture angle is larger at higher frequencies. The posture angle in the previous studies was assuming a normal distribution with a mean of -5° and a standard deviation of 15° ; however, the mean swimming angle in this study was -1.23° with a standard deviation of 11.5° . Since the chub mackerel tended to turn their bodies downward by approximately 3° relative to the direction of travel (Nauen and Lauder 2002), assuming this 3° relative to the swimming angle in this study, the mean posture angle is -4.23° , which is 0.77° more parallel to the previous study. Because the directivity of TS with respect to posture angle becomes

Table 4 Coefficients of $TS_{\text{mean}}^{\text{fork length (FL)}}$, $TS_{\text{mean}}^{\text{body length (L)}}$ and $TS_{\text{mean}}^{\text{total length (TL)}}$ estimates by previous studies for chub mackerel and their related species plotted in Fig. 7

Reference	Method	Species	Size range (cm)	Frequency (kHz)	TS_{cm}
This study	Free-swimming	<i>Scomber Japonicus</i>	FL: 17.4–34.0	38	–67.9
				120	–69.2
Kuznetsov et al	In situ		L: 20.0–30.0	38	–72.8
				120	–73.6
Park et al	Ex situ		FL: 15.4–26.2	38	–66.0
				120	–66.0
Lee et al	Ex situ		L: 26.2–38.3	50	–67.2
				120	–66.9
Tong et al	Ex situ		L: 12.0–22.2	38	–73.3
				120	–74.2
Gutierrez et al	In situ	<i>Scomber scombrus</i>	L: 26.0–30.0	38	–70.9
				120	–70.8
Scoulding et al	In situ		TL: Ave. 33.3	38	–90.1
				120	–86.7
Palermino et al	In situ		L: 31.6–40.6	38	–71.6
				120	–74.8
Kawauchi et al	Free-swimming	<i>Trachurus japonicus</i>	FL: 12.5–27.5	38	–66.7
				120	–69.5

stronger at high frequencies, it is possible that this slight difference is more significant at 120 kHz than at 38 kHz, and that the TS_{cm} at 120 kHz in this study was estimated lower.

Additionally, the TS_{cm} of chub mackerel (12.04–22.17 cm) estimated by the Kirchhoff-ray mode (KRM) model, which uses average posture angle N (-3° , 4°), was -73.27 dB at 38 kHz and -74.18 dB at 120 kHz, which were 3.2–5.3 dB weaker than the estimates in the present study (Tong et al. 2022). This can be explained by the difference between the parameters used in the theoretical model and those relevant to natural conditions. For example, theoretical values were used for the probability density distribution of posture angle and the ratio g between the fish body and seawater (different acoustic impedances), which are important parameters for model calculations (Matsukura et al. 2013). For theoretical values, g is assumed to be uniform within the fish body; however, there are multiple boundaries even within the fish body. Since the g is affected by the posture angle, it is quite difficult to predict the movement of fish bodies under natural conditions and to account for these changes in the calculation of TS . Therefore, it is considered that the assumed parameters in the theoretical cannot reflect all the conditions of the fish in the natural environment and that this may have led to a low TS_{cm} estimate. We believe that the results of this study are more realistic than those of previous studies because they account for the swimming characteristics of chub mackerel.

Finally, the equation for the TS – FL relationship in this study was compared with those for closely related species and other small-pelagic fish (Table 4, Fig. 7). First, we

compared our TS values with those of the Atlantic mackerel, which is closely related to chub mackerel. Many estimates obtained by in situ methods were significantly lower than those in the present study, even for the same length class (Palermino et al. 2021; Gutierrez and MacLennan 1998; Scoulding et al. 2017). The low TS in the Atlantic mackerel can be explained by its lack of a swim bladder. The difference in TS due to the influence of the swim bladder has been verified using a theoretical scattering model; even for fish of the same shape, the TS of bladder-less fish was more than 10 dB lower than that of fish with a swim bladder (Sawada 2002). On the other hand, the TS_{cm} of Japanese jack mackerel, which has a similar shaped physoclistous swim bladder to that of chub mackerel, was estimated using the same method as in this study, yielding similar results (i.e., -66.7 dB at 38 kHz and -69.5 dB at 120 kHz) (Kawauchi et al. 2019). In the future, we expect the method used in this study to be useful for obtaining accurate TS values for other fish species under natural conditions.

Future prospects

In this study, we collected free-swimming chub mackerel TS data over a wide length range, enabling us to derive highly accurate TS data (i.e., close to values in natural conditions) and to derive a highly accurate TS –fork length relationship. However, our analysis did not fully account for the posture angle and swimming speed of school swimmers. For fish species that swim in schools in the wild, it is necessary to obtain measurements in schooling conditions.




Fig. 7 Estimated TS –length relationship of chub mackerel and other species from the literature and this study. *S. S.* is an abbreviation for Atlantic mackerel and *T. J.* is an abbreviation for Japanese jack mackerel. Kuznetsov et al. (38 kHz: $20\log L-72.8$; 120 kHz: $20\log L-73.6$), Park et al. (38 kHz: $20\log FL-66.0$; 120 kHz: $20\log FL-66.0$), Lee et al. (50 kHz: $20\log L-67.2$; 120 kHz: $20\log L-66.9$), Tong et al. (38 kHz: $20\log L-73.3$; 120 kHz: $20\log L-74.2$), Gutierrez et al. (38 kHz: $20\log L-70.9$; 120 kHz: $20\log L-70.8$), Scouling et al. (38 kHz: $20\log TL-90.1$; 120 kHz: $20\log TL-86.7$), Palermino et al. (38 kHz: $20\log L-71.6$; 120 kHz: $20\log L-74.8$), Kawauchi et al. (38 kHz: $20\log FL-66.7$; 120 kHz: $20\log FL-69.5$)

Therefore, further studies are needed to improve the accuracy of the TS –fork length relationship, such as analyses of TS in schools.

The frequencies used in this study are carried by many research vessels, and we expect the TS –fork length relationship obtained in this study to be used for high-precision estimation of the abundance of chub mackerel in the future. Additionally, the TS_{mean} at 38 kHz was slightly higher than that at 120 kHz for the same fork length, indicating frequency-dependent characteristics of chub mackerel. Since such differences in reflection intensity among frequencies differ among species (Kang et al. 2002; Sato et al. 2015), we believe that these frequency characteristics can be used for species identification in the sea. In the future, our approach can be applied to measure TS in more fish species and to clarify practical TS and frequency characteristics. Our results are expected to provide a basis for the estimation of the abundance of fish species in different areas with a high degree of accuracy.

Acknowledgements We would like to express our deepest appreciation to Marine Fisheries Research and Development Center, Japan Fisheries Research and Education Agency for supporting our study. We thank all members of the Laboratory of Marine Ecosystem Change Analysis for their support. We also thank Editage (www.editage.com) for English language editing.

References

- Abe K, Iida K, Mukai T (1999) Diurnal changes of area backscattering coefficient during the acoustic survey of Walleye Pollock. *Nippon Suisan Gakkaishi* 65(2):252–259. <https://doi.org/10.2331/suisan.65.252>
- Amakasu K, Furusawa M, Fan C (2003) A method for target strength measurement of small-size animals by split-beam method in small tank. *Nippon Suisan Gakkaishi* 69(4):575–583. <https://doi.org/10.2331/suisan.69.575>
- Amakasu K, Sadayasu K, Abe K, Takao Y, Sawada K, Ishii K (2010) Swimbladder shape and relationship between target strength and body length of Japanese anchovy (*Engraulis japonicus*). *J Mar Acoust Soc Jpn* 37(1):46–59. <https://doi.org/10.3135/jmasj.37.46>
- Footo KG (1980) Importance of the swimbladder in acoustic scattering by fish, a comparison of gadoid and mackerel target strengths. *J Acous Soc Am* 67:2084. <https://doi.org/10.1121/1.384452>
- Furusawa M (1991) acoustic remote sensing techniques for fisheries resource surveys. *J Remote Sens Soc Jpn* 11(2):313–319. <https://doi.org/10.11440/rssj1981.11.313>
- Furusawa M, Amakasu K (2010) The analysis of echotrace was obtained by a split-beam echosounder to observe the tilt-angle dependence of fish target strength in situ. *ICES J Mar Sci* 67(2):215–230. <https://doi.org/10.1093/icesjms/fsp246>
- Furusawa M, Miyanoohana Y (1988) Application of echo-trace analysis to estimation of behavior and target strength of fish. *J Acous Soc Jpn* (e) 9(4):169–180. <https://doi.org/10.1250/ast.9.169>
- Furusawa M, Suzuki H, Miyanoohana Y (1989) A multipurpose quantitative echo sounding system. *Nat Res Instit Fish Eng* 16(2):82–93. <https://doi.org/10.11489/jmasj1986.16.82>
- Gibb AC, Dickson KA, Lauder GV (1999) Tail kinematics of the chub mackerel *Scomber japonicus*, testing the homocercal tail model of fish propulsion. *J Exp Biol* 202(18):2433–2447. <https://doi.org/10.1242/jeb.202.18.2433>
- Goto N, Hashida D, Nakao H, Tomiyama T (2020) Spatiotemporal distribution of two mackerel species, *Scomber japonicus* and *Scomber australasicus*, in the Bungo Channel, western Japan. *Nippon Suisan Gakkaishi* 86(4):295–301. <https://doi.org/10.2331/suisan.19-00043>
- Gutierrez M, MacLennan D N (1998) Resultados preliminares de las mediciones de fuerza de blanco in situ de las principales especies pelágicas. *Crucero BIC Humboldt 9803–05 de Tumbes a Tacna*. Instituto del Mar del Perú—IMARPE 135: 16–19. (in Spanish with English abstract)
- Hasegawa E, Kobayashi H (1989) Estimate of the trace of swimming behavior of fish around underwater lamps. *Nippon Suisan Gakkaishi* 55(10):1707–1714. <https://doi.org/10.2331/suisan.55.1707>
- Henderson MJ, Horne JK, Towler RH (2008) The influence of beam position and swimming direction on fish target strength. *ICES J Mar Sci* 65(2):226–237. <https://doi.org/10.1093/icesjms/fsm190>
- Hirose M, Mukai T, Hwang D, Iida K (2005) Target strength measurements on tethered live jellyfish *Nemopilema nomurai*. *Nippon Suisan Gakkaishi* 71(4):571–577. <https://doi.org/10.2331/suisan.71.571>
- Ichinokawa M, Okamura H (2014) Review of stock evaluation methods using VPA for fishery stocks in Japan, implementation with R. *Bull Jpn Soc Fish Oceanogr* 78(2):104–113

- Ichinokawa M, Okamura H, Kurota H (2017) The status of Japanese fisheries relative to fisheries around the world. *ICES J Mar Sci* 74(5):1277–1287. <https://doi.org/10.1093/icesjms/fsx002>
- Ito M, Matsuo I, Imaizumi T, Akamatsu T, Wang Y, Nishimori Y (2015) Target strength spectra of tracked individual fish in schools. *Fish Sci* 81(4):621–633. <https://doi.org/10.1007/s12562-015-0890-7>
- Kanehiro H, Suzuki M, Matuda K (1985) Characteristics of schooling behavior by the group size of rose bitterling in the experimental water tank. *Bull Jpn Soc Sci Fish* 51(12):1977–1982. <https://doi.org/10.2331/suisan.51.1977>
- Kang M, Furusawa M, Miyashita K (2002) Effective and accurate use of difference in mean volume backscattering strength to identify fish and plankton. *ICES J Mar Sci* 59(4):794–804. <https://doi.org/10.1006/jmsc.2002.1229>
- Kawai H, Yatsu A, Watanabe C, Mitani T, Katsukawa T, Matsuda H (2002) Recovery policy for chub mackerel stock using recruitment-per-spawning. *Fish Sci* 68:963–971. <https://doi.org/10.1046/j.1444-2906.2002.00520.x>
- Kawauchi Y, Minami K, Shirakawa H, Miyashita K, Iwahara Y, Tomiyasu M, Kobayashi M, Sakai T, Shao H, Nakagawa M (2019) Target strength measurement of free-swimming jack mackerel using an indoor large experiment tank. *Nippon Suisan Gakkaishi* 85(1):2–16. <https://doi.org/10.2331/suisan.18-00008>
- Kurota H, Kitajima S, Goto T, Sassa C, Tanaka S, Hiramatsu K, Muto S, Yasuda T, Yamada A, Yamada H, Yukami R, Yoda M (2019) Fisheries biology and population dynamics of Tsushima Warm Current stock of chub mackerel *Scomber japonicas*. *Bull Jpn Soc Fish Oceanogr* 84(4):237–251
- Kuznetsov MY, Polyanchko VI, Syrovatkin EV, Ubarchuk IA (2021) The study of the acoustic target strength of Chub mackerel in the northwestern Pacific Ocean in situ. *Equip Fish Res* 184:73–86. <https://doi.org/10.36038/2307-3497-2021-184-73-86>. ((in Russian with English abstract))
- Lee D, Shin H (2005) Construction of a data bank for acoustic target strength with fish species, length and acoustic frequency for measuring fish size distribution. *J Korean Soc Fish Ocean Technol* 38(4):265–275. <https://doi.org/10.5657/kfas.2005.38.4.265>
- Matsukura R, Sawada K, Abe K, Minami K, Nagashima H, Yonezaki S, Murase H, Miyashita K (2013) Comparison of measurements and model calculations of target strength of juvenile sandeel in Sendai Bay. *Nippon Suisan Gakkaishi* 79(4):638–648. <https://doi.org/10.2331/suisan.79.638>
- Medwin H, Clay CS (1988) Fundamentals of acoustical oceanography (in applications of modern acoustics). Acad Press San Diego. <https://doi.org/10.1016/B978-0-12-487570-8.X5000-4>
- Mitson RB (1983) Fisheries sonar. Fishing News Books, England
- Miyanohana Y (1994) Study on the characteristics of the target strength of fish. *Bull Nat Res Instit Fish Eng* 15:83–144
- Miyanohana Y, Ishii K, Furusawa M (1990) Dorsal aspect target strength functions of eight species of fish at four frequencies. *Tech Rep Nat Res Instit Fish Eng* 10:65–154
- Miyashita K (2019) Development and application of acoustic methods for monitoring fisheries resources and ecosystems. *Nippon Suisan Gakkaishi* 85(3):294–296. <https://doi.org/10.2331/suisan.WA2632>
- Mukai T, Sano N, Iida K, Sasaki S (1993) Studies on dorsal aspect target strength of ten species of fish collected in the East China Sea. *Nippon Suisan Gakkaishi* 59(9):1515–1525. <https://doi.org/10.2331/suisan.59.1515>
- Mukai T, Sano N, Iida K, Sasaki S (1994) Relationship between dorsal aspect target strength of fish caught in the East China Sea and their Swimbladder. *Nippon Suisan Gakkaishi* 60(2):215–222. <https://doi.org/10.2331/suisan.60.215>
- Muko S, Kurota H (2018) Comparison of stock assessments based on calendar year and fishing year, an example of the Tsushima Warm Current stock of chub mackerel. *Bull Jpn Soc Fish Oceanogr* 82(1):14–25. https://doi.org/10.34423/jso.82.1_14
- Munoz L, Aspillaga E, Palmer M, Saraiva JL, Arechavala-Lopez P (2020) Acoustic telemetry, a tool to monitor fish swimming behavior in sea-cage aquaculture. *Front Mar Sci* 7(645):1–12. <https://doi.org/10.3389/fmars.2020.00645>
- Nakayama T, Hiramatsu K (2010) Evaluation of the reliability of VPA used for stock assessment for TAC species. *Nippon Suisan Gakkaishi* 76(6):1043–1047. <https://doi.org/10.2331/suisan.76.1043>
- Nauen JC, Lauder GV (2002) Hydrodynamics of caudal fin locomotion by chub mackerel, *Scomber japonicas* (Scombridae). *J Exp Biol* 205(12):1709–1724. <https://doi.org/10.1242/jeb.205.12.1709>
- Okada M (2020) Changes in the resource biological characteristics of chub mackerel *Scomber japonicas* in the Kumano-nada due to changes in stock levels. *Fish Biol Oceanogr Kuroshio* 21:107–121
- Okunishi T, Yokouchi K, Hasegawa D, Tanaka T, Setou T, Yukami R, Kuroda H, Watanabe K, Takasuka A (2020) Relationship between sea temperature variation and fishing ground formations of chub mackerel in the Pacific Ocean off Tohoku. *Bull Jpn Soc Fish Oceanogr* 84(4):271–284. https://doi.org/10.34423/jso.84.4_271
- Palermينو A, Felice AD, Canduci G, Biagiotti I, Costantini I, Malavolti S, Leonori I (2021) First target strength measurement of *Trachurus mediterraneus* and *Scomber colias* in the Mediterranean Sea. *Fish Res* 240:105973. <https://doi.org/10.1016/j.fishres.2021.105973>
- Park G, Oh W, Oh S, Lee K (2022) Acoustic scattering characteristics of chub mackerel (*Scomber japonicus*) by KRM model. *J Korean Soc Fish Ocean Technol* 58(1):32–38. <https://doi.org/10.3796/KSFOT.2022.58.1.032>. (in Korean with English abstract)
- Sassa C, Tsukamoto Y (2010) Distribution and growth of *Scomber japonicus* and *S. australasicus* larvae in the southern East China Sea in response to oceanographic conditions. *Mar Ecol Prog Ser* 419:185–199. <https://doi.org/10.3354/meps08832>
- Sato M, Horne JK, Parker-Stetter SL, Keister JE (2015) Acoustic classification of coexisting taxa in a coastal ecosystem. *Fish Res* 172:130–136. <https://doi.org/10.1016/j.fishres.2015.06.019>
- Sawada K (2002) Study on the precise estimation of the target strength of fish. *Bull Fish Res Agency* 2:47–122
- Scoulling B, Gastauer S, MacLennan DN, Fassler SM, Coplang P, Fernandes PG (2017) Effects of variable mean target strength on estimates of abundance, the case of Atlantic mackerel (*Scomber scombrus*). *ICES J Mar Sci* 74(3):822–831. <https://doi.org/10.1093/icesjms/fsw212>
- Shimura T, Ohshimo S, Terakado H, Tian Y (2009) Development of method for estimation of 0+ fish abundance index of jack mackerel *Trachurus japonicus* by mid-water trawling and area-swept method in the southwestern Sea of Japan. *Nippon Suisan Gakkaishi* 75(6):1042–1050. <https://doi.org/10.2331/suisan.75.1042>
- Simmonds J, MacLennan DN (2005) Fisheries acoustics-theory and practice, 2nd edn. Blackwell Publishing, Oxford
- Simmonds EJ, Williamson NJ, Gerlotto F, Aglen A (1992) Acoustic survey design and analysis procedure, a comprehensive review of current practice. *ICES Cooperative Res Rep* 187:94–95
- Tanigami S, Fukuda M, Mukai T, Iida K (2006) Target strength estimation of the Japanese anchovy *Engraulis japonicus* using control methods and acoustic scattering models. *Jpn Soc Fish Eng*. https://doi.org/10.18903/pamjsfe.2006.0_295
- Tong J, Xue M, Zhu Z, Wang W, Tian S (2022) Impacts of morphological characteristics on target strength of Chun Mackerel (*Scomber japonicus*) in the Northwest Pacific Ocean. *Front Mar Sci* 9:856483. <https://doi.org/10.3389/fmars.2022.856483>
- Wardle C S (1980) NATO Advanced Study Institutes Series, Environmental Physiology of Fishes, Effects of Temperature on the Maximum Swimming Speed of Fishes. Springer, Boston.
- Yatsu A, Kaeriyama M (2005) Linkages between coastal and open-ocean habitats and dynamics of Japanese stocks of chum salmon

- and Japanese sardine. *Deep-Sea Res II* 52:727–737. <https://doi.org/10.1016/j.dsr2.2004.12.019>
- Yatsu A, Takahashi K, Watanabe K, Honda O (2019) Seasonal and interannual variability in crude fat content and condition factor of chub mackerel *Scomber japonicus* captured from waters off northeastern Japan during 2012–2017. *Bull Jpn Soc Fish Oceanogr* 83(1):19–27
- Yilmaz Ö (1987) Seismic data processing. Society of Exploration Geophysicists, Tulsa
- Zhu Y, Minami K, Iwahara Y, Oda K, Hidaka K, Hoson O, Morishita K, Hirota M, Tsuru S, Shirakawa H, Miyashita K (2021a) Seasonal variation in fish school spatial distribution and abundance under the Kuroshio regular pattern and the large meander in Suzu coastal waters. *PLoS ONE* 16(11):e0260629. <https://doi.org/10.1371/journal.pone.0260629>
- Zhu Y, Minami K, Iwahara Y, Oda K, Hidaka K, Hoson O, Morishita K, Hirota M, Tsuru S, Shirakawa H, Miyashita K (2021b) Seasonal dynamics in fish distribution and abundance revealed by an acoustic survey in coastal waters of the Suzu Area, Kochi Prefecture, Japan. *J Mar Sci Technol* 29(2):146–157. <https://doi.org/10.51400/2709-6998.1086>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.